

Juvenile Myelomonocytic Leukemia (JMML) With the Hematologic Phenotype of Severe β Thalassemia

George R. Honig,^{1*} Carlos R. Suarez,² Loyda N. Vida,¹ Shi-Jiang Lu,¹ and Edison T. Liu³

¹Department of Pediatrics, University of Illinois College of Medicine, Chicago

²Department of Pediatrics, Loyola University Medical Center, Maywood, Illinois

³Lineberger Cancer Center, University of North Carolina, Chapel Hill

A 3-year-old Filipino-American child with recurrent fever, splenomegaly, anemia, and thrombocytopenia, was found to have a hemoglobin F level of 76.9%. His reticulocyte count was elevated (4.3%), and erythroblasts were present in his peripheral blood. The child's erythrocytes were microcytic (MCV 66.9 fl) but his serum ferritin level was normal. His bone marrow at initial presentation demonstrated normal cellularity without an increase in blast cells. The disease progressed with worsening anemia, leukocytosis, and thrombocytopenia, with increased blasts in his marrow and the appearance of a mediastinal mass. His liver, spleen, and lymph nodes were found to be infiltrated with myeloblasts, supporting a diagnosis of juvenile myelomonocytic leukemia (JMML). Analysis of the child's Hb F showed a G_{γ}/A_{γ} ratio of 2.2, which was within the characteristic range for JMML. A globin synthesis study using blood reticulocytes showed an $\alpha/\text{non-}\alpha$ globin synthesis ratio of 2.24, typical of severe homozygous β thalassemia. Southern blot analysis of blood-leukocyte DNA from the patient and his parents demonstrated no apparent abnormality in the β -globin gene promoter or coding regions. The elevated level of Hb F in this child with JMML appeared to be part of an acquired Cooley's anemia-like hematologic phenotype. *Am. J. Hematol.* 58:67–71, 1998. © 1998 Wiley-Liss, Inc.

Key words: juvenile myelomonocytic leukemia; fetal hemoglobin; hemoglobin synthesis

INTRODUCTION

Juvenile myelomonocytic leukemia (JMML), previously referred to as juvenile chronic myeloid leukemia, is an uncommon disease of infants and young children. The disorder is expressed as a clonal panmyelopathy [1–3] that is distinct from Philadelphia chromosome-positive adult CML. Children with type I neurofibromatosis account for 7% of the reported cases of JMML [1,4]. Approximately 30% of JMML patients have been shown to have mutations of the *N-ras* proto-oncogene [5].

Typical clinical features of JMML include leukocytosis with the presence of early myeloid and monocytic elements, thrombocytopenia, skin rash, and hepatosplenomegaly. Progression of the disease is often rapid, with infiltration of the bone marrow and other tissues with monocytic and myeloid cells.

A remarkable finding in a majority of children with the JMML syndrome is the presence of strikingly elevated levels of fetal hemoglobin (Hb F) [6]. The G_{γ}/A_{γ} ratio of the Hb F from these patients has been found to be similar to that of Hb F from normal newborn infants [7–9], which, taken together with their characteristically low

levels of carbonic anhydrase and high i antigen titers, has suggested that erythropoiesis in JMML may reflect a reversion to a fetal pattern. Studies of Hb F synthesis by BFU-E-derived bursts [8] have provided further support for this notion.

In this report, we describe the findings from a child with the JMML syndrome with greatly elevated levels of Hb F, whose pattern of hemoglobin synthesis suggested the presence of an acquired abnormality with β thalassemia-like hematologic features.

METHODS

Hemoglobin Studies

Procedures for the preparation of hemoglobins for analysis, Hb electrophoresis, and measurements of Hb F

*Correspondence to: George R. Honig, Department of Pediatrics (M/C 856), University of Illinois, 840 S. Wood St., Chicago, IL 60612. E-mail: ghonig@uic.edu

Received for publication 30 June 1997; Accepted 10 December 1997

TABLE I. Representative Hematologic Values From the Patient and His Parents

	Patient	Mother	Father
Hemoglobin (G/dl)	10.5	14.8	14.9
MCV (fl)	67	89	95
Reticulocytes (%)	4.3	0.5	0.8
Platelets ($\times 10^{-3}$)	32	255	214
Hemoglobin electrophoresis	FA	AA	AA
Hb A ₂ (%)	0.4	2.6	3.3
Hb F (%)	76.9	0.3	0.3

and Hb A₂ were as described previously [10]. For the quantitative estimation of globin chains, HPLC was performed with a C₄ column (Vydac, The Separations Group, Hesperia, CA) with acetonitrile-water-trifluoroacetic acid gradient elution [11]. For globin synthesis studies, washed red cells were incubated for 2 hr at 37°C in medium containing L-leucine-³H; radiolabeled globin chains were fractionated by carboxymethylcellulose column chromatography [12].

DNA Analyses

High-molecular-weight DNA was prepared by phenol/chloroform extraction [13] from blood leukocytes and from formalin-fixed splenic tissue. The procedure for identification of N-ras gene mutations was as previously described [14].

For Southern blot analyses, DNA from normal controls and from the patient and his parents was digested with Bam HI, Hind III, Eco RI, or Pst I, and hybridized with a full-length human cDNA β globin-gene probe [13,15,16]. In studies that examined the 5' β globin-gene promoter region, the segment between nucleotides 40250 and 41270 [17] was amplified by PCR, followed by digestion with Bam HI.

CLINICAL FINDINGS

A previously well 3-year-old boy of Filipino parentage was evaluated because of recurrent fevers and lethargy. His liver was found to extend 1.5 cm below his costal margin, and his spleen 4 cm. No apparent rash or other abnormality was found from his examination.

His hematologic findings were: WBC 9,600/ μ l; Hb 10.5 G/dl; platelets 32,000/ μ l; RBC 4.07×10^6 / μ l; MCV 66.9 fl; and reticulocytes 4.3%. His blood counts also showed variable numbers of erythroblasts. His serum ferritin level was 78.2 μ g/ml. His bone marrow had normal cellularity with erythroid hyperplasia, but no dyserythropoietic changes or increase in blast cells. Cytogenetic analysis revealed a normal 46XY karyotype. Hemoglobin studies demonstrated a greatly elevated level of Hb F (Table I). Hematologic values from both of the parents were normal.

Over the following month, the child's spleen became progressively larger accompanied by abdominal pain. His leukocyte count increased to 60,000/ μ l, and his platelet count decreased to approximately 20,000. Examination of his bone marrow revealed an increase in blast cells to 7%. Computed tomography of his chest and abdomen demonstrated the presence of a mass bordering the right lower mediastinum, as well as periaortic lymph node enlargement. Biopsy specimens of the child's spleen, liver, and lymph nodes showed infiltration with myeloblasts, supporting a diagnosis of JMML.

A chemotherapy regimen was initiated that included dexamethasone, etoposide, daunorubicin, cytarabine, thioguanine, and 2-chlorodeoxyadenosine. The child showed no appreciable response to this treatment, and he subsequently underwent a haploidentical transplant with bone marrow from his mother. Engraftment was successfully achieved, with normalization of his hematologic values, but 5 months later he experienced a recurrence of his disease, which progressed with a rapidly downhill course.

RESULTS

Hemoglobin Findings

Electrophoresis of the patient's hemoglobin demonstrated an FA pattern with a low percentage of Hb A₂. Smears of his peripheral blood prepared according to the procedure of Kleihauer et al. [18] demonstrated strongly-positive staining of Hb F in all of the red cells. These studies from both of the parents yielded entirely normal results.

HPLC fractionation of the child's globin chains (Fig. 1) confirmed that the non- α chains were predominantly of the γ type, with a γ/α ratio of 2.2.

Globin Synthesis Results

Blood reticulocytes from the patient actively incorporated radiolabeled L-leucine into globin chains. The relative rates of synthesis of the individual globin types are shown in the chromatogram in Figure 2. The synthesis of γ chains accounted for approximately 65% of the non- α chains, in close correspondence with the percentage of Hb F in the child's blood. As the chromatogram in Figure 2 also demonstrates, α chain synthesis was substantially in excess of non- α chains, with an $\alpha/[\beta + \gamma]$ chain synthesis ratio of 2.24. Globin synthesis studies using reticulocytes from the parents yielded low levels of radioleucine incorporation into globin chains; the α /non- α globin synthesis ratios from both of them were within the range of normal.




Fig. 1. HPLC fractionation showing the representation of globin chains in the patient's hemoglobin.




Fig. 2. Incorporation of L-leucine-³H into globin chains by reticulocytes from the patient's blood.

DNA Studies

DNA from the patient's blood leukocytes and splenic tissue showed no evidence of mutation in the *N-ras* gene. Southern blot analyses of the β globin gene coding region and upstream promoter region, using DNA from the patient and his parents, yielded restriction enzyme frag-

ments that invariably were indistinguishable from those of normal controls (data not shown).

HLA/DR Studies

Histocompatibility testing of the patient showed him to be: A11, A34, B40, B62, DR 12, DR 15. Results from his mother were: A24, A34, B35, B40, DR 12, DR 15; and from his father: A11, A34, B62, DR4, DR 12.

DISCUSSION

The nonspecific clinical findings early in the course of this child's illness made it difficult initially to arrive at a specific diagnosis. As his disease progressed, with the appearance of blast cells in his bone marrow and in other tissues, and in the face of his characteristic pattern of Hb F expression, it became apparent that this syndrome represented JMML. Although elevated levels of Hb F as an acquired abnormality have been identified in patients with a variety of forms of leukemia, Hb F levels in excess of 50% appear to be unique to the JMML syndrome [19].

The increased $G\gamma/A\gamma$ ratio of this patient's Hb F, together with his very low level of Hb A₂, were entirely comparable to findings from other children with JMML [7,8,9,19]. These specific changes, however, in addition to occurring in the fetus and newborn, have also been observed in patients with severe forms of β thalassemia from a wide variety of ethnic backgrounds [20–24].

Our findings from the globin synthesis study of this patient (shown in Fig. 2) included a highly unbalanced synthesis pattern, with a substantial excess of α chain synthesis in relation to that of the non- α chains. These results were indistinguishable from those obtained in similar studies of patients with severe β thalassemia [25,26].

This child's clinical syndrome also included many of the other characteristic hematologic features of severe β thalassemia: His blood erythrocytes were microcytic and dysmorphic, and his reticulocyte count was consistently elevated and accompanied by a significant degree of normoblastemia.

The other hematologic data from the patient and his parents provided no indication of a genetic basis for his Cooley's anemia-like syndrome: Both parents had normal hemoglobin levels as well as normocytic red cell indices, their levels of Hb A₂ and Hb F were not elevated, their erythrocytes showed no evidence of Hb F-containing cells by the Kleihauer-Betke procedure, and the patterns of globin synthesis by reticulocytes from their blood were also within the normal range. Southern blot analysis of the β -globin gene coding region and upstream promoter region from the patient's DNA also produced patterns that were indistinguishable from those of normal controls.

We are not aware of other studies of globin synthesis from patients with JMML, and it may be possible that the elevated levels of Hb F in others of these patients also represent part of an acquired β thalassemia-like syndrome.

Hematologic disorders having many features in common with α thalassemia, expressed with the phenotype of Hb H disease, have been observed as acquired abnormalities in patients with a wide variety of other blood diseases, including acute lymphoblastic leukemia [27], erythroleukemia [28,29], adult CML [30], sideroblastic anemia [31,32], myelofibrosis [33,34], and other forms of myeloproliferative disease [35,36].

Efforts to identify the underlying molecular abnormality in these acquired Hb H disease syndromes have been only partly successful. Studies of several of these patients [32,36,37] have shown markedly decreased levels of α -chain mRNA in their hemoglobin-synthesizing cells. However restriction enzyme mapping studies of their α -globin and ζ -globin genes were unrevealing of any abnormality [36,37].

In an effort to determine if the decreased α -globin gene expression in these patients was directly related to the α -globin genes on chromosome 16, Helder and Deisseroth [38] performed experiments in which the chromosome 16 from several affected individuals was transferred to mouse erythroleukemia cells. S1 nuclease analysis showed that in a number of such mouse cell lines, human α -globin gene expression was similar to that of cells containing human chromosome 16 obtained from normal individuals. These observations suggested, therefore, that the suppression of α -globin synthesis in these patients with the acquired Hb H syndrome might well be the result of a genetic abnormality involving a factor acting in trans to the α -globin gene.

Acquired forms of microcytic anemia with β thalassemia-like expression have been described only rarely. In reports of two such syndromes [39,40], one involving an adult with erythroleukemia and the other an elderly individual with refractory anemia accompanied by ring sideroblasts, both of the patients were described as having high levels of Hb F with lower-than-normal levels of Hb A₂; In light of these findings, both of these acquired syndromes were characterized as being $\delta\beta$ thalassemia-like.

Globin synthesis studies of additional patients with JMML will be needed to test our hypothesis that the elevated level of Hb F associated with this disease may be one element of an acquired β thalassemia-like syndrome. If this indeed proves to be the case, JMML patients could provide a further opportunity to gain an understanding of the underlying molecular basis of these acquired abnormalities.

ACKNOWLEDGMENTS

We thank Ms. Julia Ferrari for her expert technical assistance.

REFERENCES

1. Aricò M, Biondi A, Pui C-H: Juvenile myelomonocytic leukemia. *Blood* 90:479–488, 1997.
2. Amenomori T, Tomonaga M, Yoshida Y, Kuriyama K, Matsuo T, Jinnai L, Ichimaru M, Omiya A, Tsuji Y: Cytogenetic evidence for partially committed myeloid progenitor cell origin of chronic myelomonocytic leukaemia and juvenile chronic myeloid leukaemia: Both granulocyte-macrophage precursors and erythroid precursors carry identical marker chromosome. *Br J Haem* 64:539–546, 1986.
3. Busque L, Gilliland DG, Prchal JT, Sieff CA, Weinstein HJ, Sokol JM, Belickova M, Wayne AS, Zuckerman KS, Sokol L, Castleberry RP, Emanuel PD: Clonality in juvenile chronic myelogenous leukemia. *Blood* 85:21–30, 1995.
4. Shannon KM, O'Connell P, Martin GA, Paderanga D, Olson K, Dinndorf P, McCormick F: Loss of the normal *NF1* allele from the bone marrow of children with type I neurofibromatosis and malignant myeloid disorders. *N Engl J Med* 330:597–601, 1994.
5. Miyauchi J, Asada M, Sasaki M, Tsunematsu Y, Kojima S, Mizutani S: Mutations of the *N-ras* gene in juvenile chronic myelogenous leukemia. *Blood* 83:2248–2254, 1994.
6. Weatherall DJ, Edwards JA, Donohoe WTA: Haemoglobin and red cell enzyme changes in juvenile myeloid leukaemia. *Br Med J* 1:679–681, 1968.
7. Maurer HS, Vida LN, Honig GR: Similarities of the erythrocytes in juvenile chronic myelogenous leukemia to fetal erythrocytes. *Blood* 39:778–784, 1972.
8. Weinberg RS, Leibowitz D, Weinblatt ME, Kochen J, Alter BP: Juvenile chronic myelogenous leukaemia: The only example of truly fetal (not fetal-like) erythropoiesis. *Br J Haematol* 76:307–310, 1990.
9. Huisman THJ, Schroeder WA: Fetal hemoglobin synthesis in some forms of leukemia. In Peeters H (ed): "Protides of Biological Fluids, Proceedings of the 24th Colloquium," Volume 24. Oxford: Pergamon Press, 1976.
10. Honig GR, Shamsuddin M, Vida LN, Mompont M, Valcourt E, Bowie LJ, Jones EC, Powers PA, Spritz RA, Guis M, Embury SH, Conboy J, Kan YW, Mentzer WC, Weil SC, Hirata RK, Waloch J, O'Riordan JF, Goldstick TK: Hemoglobin Evanston (α 14 Trp \rightarrow Arg): An α chain variant expressed as α thalassemia. *J Clin Invest* 73:1740–1749, 1984.
11. Strahler JR, Rosenblum BB, Hanash SM: A silent, neutral substitution detected by reverse-phase high-performance liquid chromatography: Hemoglobin Beirut. *Science* 221:860–862, 1983.
12. Honig GR, Mason RG, Vida LN, Shamsuddin M: Synthesis of hemoglobin Abraham Lincoln (β 32 Leu \rightarrow Pro). *Blood* 43:657–664, 1974.
13. Lu S-J, Rowan S, Bani MR, Ben-David Y: Retroviral integration within the Fli-2 locus results in inactivation of the erythroid transcription factor NF-E2 in Friend erythroleukemia: Evidence that NF-E2 is essential for globin expression. *Proc Natl Acad Sci USA* 91:8398–8402, 1994.
14. Neubauer A, Shannon S, Liu E: Mutations of the *ras* proto-oncogenes in childhood monosomy 7. *Blood* 77:594–598, 1991.
15. Ben-David Y, Giddens E, Bernstein A: Identification and mapping of a common proviral integration site *Fli-1* in erythroleukemic cells induced by Friend murine leukemia virus. *Proc Natl Acad Sci USA* 87:1332–1336, 1990.
16. Feinberg AP, Vogelstein B: A technique for radiolabeling DNA restriction fragments to high specific activity. *Anal Biochem* 132:6–13, 1983.

17. Collins FS, Weissman SM: The molecular genetics of human hemoglobins. *Prog Nucleic Acid Res Mol Biol* 31:315–462, 1984.
18. Kleihauer E, Braun H, Betke K: Demonstration von fetalem Hämoglobin in die Erythrocyten eines Blutaussstrichs. *Klin Wochenschr* 35: 637–638, 1957.
19. Weatherall DJ, Pembrey ME, Pritchard J: Fetal haemoglobin. *Clinics Haematol* 3:467–508, 1974.
20. Schroeder WA, Huisman THJ, Shelton JR, Shelton JB, Apell G, Bouver N: Heterogeneity of fetal hemoglobin in β -thalassemia of the negro. *Am J Hum Genet* 22:505–514, 1970.
21. Rachmilewitz EA, Huisman THJ, Schroeder WA: Heterogeneity of fetal hemoglobin among Israel families with β -thalassemia. *Israel J Med Sci* 9:1464–1468, 1973.
22. Huisman THJ, Schroeder WA, Brown AK, Hyman CB, Ortega JA, Sukumaran PK: Further studies of the postnatal change in chemical heterogeneity of human fetal hemoglobin in several abnormal conditions. *Pediatr Res* 9:1–6, 1975.
23. Lie-Injo LE, Randhawa ZI, Kane JP, Ganesan J, George R: Heterogeneity of hemoglobin gamma chains in normal newborns and in cases of alpha and beta thalassemia. *Am J Hematol* 6:17–25, 1979.
24. Comi P, Giglioni B, Ottolenghi S, Gianni AM, Ricco G, Mazza U, Saglio G, Camaschella C, Pich PG, Gianazza E, Righetti PG: γ and $\alpha\gamma$ globin chains separation and quantitation by isoelectric focusing. *Biochem Biophys Res Commun* 87:1–8, 1979.
25. Bargellesi A, Pontremoli S, Menini C, Conconi F: Excess of α -globin synthesis in homozygous β -thalassemia and its removal from the red blood cell cytoplasm. *Eur J Biochem* 3:364–368, 1968.
26. Weatherall DJ, Clegg JB, Na-Nakorn S, Wasi P: The pattern of disordered haemoglobin synthesis in homozygous and heterozygous β -thalassaemia. *Br J Haematol* 16:251–267, 1969.
27. Kueh YK: Acute lymphoblastic leukemia with brilliant cresyl blue erythrocytic inclusions: Acquired hemoglobin H? *N Engl J Med* 307: 193–194, 1982.
28. Rosenzweig AI, Heywood JD, Motulsky AG, Finch CA: Hemoglobin H as an acquired defect of alpha-chain synthesis. *Acta Haematol* 38: 91–101, 1968.
29. Pagnier J, Labie D, Kaplan JC, Junien C, Najman A, Leroux JP: Étude biochimique d'un cas d'érythroleucémie. *Nouv Rev Fr Hematol* 12: 317–332, 1972.
30. Beaven GH, Stevens BL, Dance N, White JC: Occurrence of haemoglobin H in leukaemia. *Nature* 199:1297–1298, 1963.
31. Boehme WM, Piira TA, Kurnick JE, Bethlenfalvay NC: Acquired hemoglobin H in refractory sideroblastic anemia. A preleukemic marker. *Arch Intern Med* 138:603–606, 1978.
32. Yoo D, Schechter GP, Amigable AN, Nienhuis AW: Myeloproliferative syndrome with sideroblastic anemia and acquired hemoglobin H disease. *Cancer* 45:78–83, 1980.
33. Dash S, Dash RJ: Idiopathic myelofibrosis without splenomegaly and with an acquired haemoglobin disorder. *Indian J Cancer* 17:193–195, 1980.
34. Veer A, Kosciolk BA, Bauman AW, Rowley PT: Acquired hemoglobin H disease in idiopathic myelofibrosis. *Am J Hematol* 6:199–206, 1979.
35. Hamilton RW, Schwartz E, Atwater J, Erslev AJ: Acquired Hb H disease. *N Engl J Med* 285:1217–1221, 1971.
36. Higgs DR, Wood WG, Barton C, Weatherall DJ: Clinical features and molecular analysis of acquired hemoglobin H disease. *Am J Med* 75:181–191, 1983.
37. Anagnou NP, Ley TJ, Chesbro B, Wright G, Kitchens C, Liebhaver S, Nienhuis AW, Deisseroth AB: Acquired α -thalassemia in preleukemia is due to decreased expression of all four α -globin genes. *Proc Natl Acad Sci USA* 80:6051–6055, 1983.
38. Helder J, Deisseroth A: S1 nuclease analysis of α -globin gene expression in preleukemic patients with acquired hemoglobin H disease after transfer to mouse erythroleukemia cells. *Proc Natl Acad Sci USA* 84:2387–2390, 1987.
39. Markham RE, Butler F, Goh K, Rowley P: Erythroleukemia manifesting $\delta\beta$ -thalassemia. *Hemoglobin* 7:71–78, 1983.
40. Aoyagi Y, Akimoto H, Yamaoka K, Koike A, Kanayama M, Miyaji T, Yamamoto K: Refractory anemia with ringed sideroblasts complicated with $\delta\beta$ -thalassemia-like hemoglobinopathy. *Rinsho Ketsueki* 30:674–679, 1989.